

IRE1 α (PT0365R) PT[®] Rabbit mAb

CatalogNo: YM8218 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat,

Applications

- WB, IHC, IF, IP, ELISA

MW

- 110kD (Calculated)
110kD (Observed)

Isotype

- IgG, Kappa

Recommended Dilution Ratios

IHC 1:100-1:500

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Basic Information

Clonality Monoclonal

Clone Number PT0365R


Immunogen Information

Specificity Endogenous


Target Information

Gene name	ERN1		
Protein Name	Serine/threonine-protein kinase/endoribonuclease IRE1		
	Organism	Gene ID	UniProt ID
	Human	2081 ;	O75460 ;
	Mouse		Q9EQY0 ;
Cellular Localization	Endoplasmic reticulum membrane		
Tissue specificity	Ubiquitously expressed. High levels observed in pancreatic tissue.		
Function	Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,Function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response transcriptional activator and triggering growth arrest and apoptosis.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family.,similarity:Contains 1 KEN domain.,similarity:Contains 1 protein kinase domain.,subunit:Homodimer; disulfide-linked. Dimer formation is driven by hydrophobic interactions within the N-terminal luminal domains and stabilized by disulfide bridges. Also binds HSPA5, a negative regulator of the unfolded protein response. This interaction may disrupt homodimerization and prevent activation of ERN1.,tissue specificity:Ubiquitously expressed. High levels observed in pancreatic tissue.,		

| Validation Data



Human brain was stained with anti-IRE1α (PT0365R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-IRE1 α (PT0365R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: LnCap Lane 2: Rat Prostate Predicted band size: 110kDa
Observed band size: 110kDa

Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 408-747-0189 (USA) 400-8787-807(China)
 Website: <http://www.immunoway.com>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
IRE1 α (PT0365R)
PT® Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)